

Digital Orthopedics and Smart Rehabilitation: The Expanding Role of Wearable Technologies and Motion Analytics in Musculoskeletal Healthcare

Jacqueline Dorian López Molina
Universidad San Carlos de Guatemala
jacquelinelopezmolina@gmail.com
<https://orcid.org/0009-0003-9030-4373>

Milagros Bartolini
Independiente
milagrosbatolini18@gmail.com
<https://orcid.org/0009-0004-7425-3198>

Óscar Javier Sepúlveda Gleiser
Complejo asistencial Dr Víctor Rios Ruiz
os.sepulveda.g@gmail.com
<https://orcid.org/0009-0006-4570-6543>

Mario René López Monzón
Universidad de San Carlos de Guatemala
marrlopezm@gmail.com
<https://orcid.org/0000-0002-2228-2227>

Juan Camilo Manosalva Trillos
Universidad de Santander
Juanamilomanosalvatrillos@gmail.com
<https://orcid.org/0009-0001-5005-9292>

Pedro Raúl Abad Valencia
Universidad de Guayaquil
pedro.abadv@ug.edu.ec
<https://orcid.org/0009-0000-5158-8949>

José Gabriel Benalcázar Game
Universidad de Guayaquil
josé.benalcazarg@ug.edu.ec
<https://orcid.org/0009-0008-9824-8480>

Hugo Eduardo Jara Sánchez
Universidad Católica de Cuenca
hugosedu@hotmail.com
<https://orcid.org/0000-0002-4021-0702>

Received: 23-Jun-2026 | Accepted: 23-Jun-2026 | Published: 28-Jun-2026

* Corresponding Author: jacquelinelopezmolina@gmail.com

How to cite this article: López Molina, J. D., Bartolini, M., Sepúlveda Gleiser, Ó. J., López Monzón, M. R., Manosalva Trillos, J. C., Abad Valencia, P. R., Benalcázar Game, J. G., & Jara Sánchez, H. E. (2026). Digital Orthopedics and Smart Rehabilitation: The Expanding Role of Wearable Technologies and Motion Analytics in Musculoskeletal Healthcare. México. International Science Journal "TheSci". 3 (1) 904-933. Quality Consulting Instituto de Educación Capacitación y Certificación de México. <https://ieccmexico.com/thesci>

Copyright (c). 2026 López Molina, J. D., Bartolini, M., Sepúlveda Gleiser, Ó. J., López Monzón, M. R., Manosalva Trillos, J. C., Abad Valencia, P. R., Benalcázar Game, J. G., & Jara Sánchez, H. E.; This is an open access article distributed under the terms of the Attribution 4.0 International (CC BY-NC-SA 4.0) International Science Journal "TheSci". Mexico Review/ Vol. 3, N. 1 / pp. 904-933/ January-June 2026 / e-ISSN: 3122-3591 / p-ISSN: 3122-3753. Research Article

ABSTRACT

Musculoskeletal disorders remain among the leading causes of disability and functional impairment worldwide, creating significant challenges for healthcare systems and rehabilitation services. Recent advances in wearable technologies, motion analytics, telemedicine, and artificial intelligence have introduced new opportunities for improving orthopedic assessment, rehabilitation monitoring, and long-term patient management. This review examines the current

applications of wearable devices in musculoskeletal healthcare, with particular emphasis on gait analysis, postoperative monitoring, rehabilitation support, remote patient management, and predictive analytics. A structured review of scientific literature indexed in major biomedical and multidisciplinary databases was conducted to evaluate the effectiveness, benefits, and limitations of wearable technologies within orthopedic practice. The findings indicate that wearable systems provide continuous and objective measurements of biomechanical and functional parameters, enabling more personalized rehabilitation strategies and supporting data-driven clinical decision-making. Gait analysis and rehabilitation monitoring emerged as the most frequently reported applications, while artificial intelligence demonstrated growing potential for predictive modeling and outcome optimization. The review also identified important implementation challenges, including data privacy concerns, interoperability limitations, device standardization issues, regulatory barriers, and patient adherence factors. Despite these challenges, wearable technologies are increasingly contributing to the development of accessible, patient-centered, and precision-based orthopedic care. The evidence suggests that continued technological innovation and integration of digital health solutions will play a fundamental role in shaping the future of musculoskeletal healthcare worldwide.

KEYWORDS

Wearable Technology, Motion Analytics, Digital Orthopedics, Musculoskeletal Disorders, Gait Analysis, Rehabilitation Monitoring, Telemedicine, Remote Patient Monitoring, Artificial Intelligence, Precision Medicine, Orthopedic Rehabilitation, Digital Health

INTRODUCTION

Musculoskeletal disorders (MSDs) represent one of the leading causes of disability, chronic pain, reduced mobility, and diminished quality of life worldwide. According to global health estimates, conditions such as osteoarthritis, low back pain, osteoporosis, sports-related injuries, and degenerative joint diseases affect hundreds of millions of individuals and impose a substantial socioeconomic burden on healthcare systems, employers, and patients [1], [2]. The growing prevalence of aging populations, sedentary lifestyles, obesity, and chronic diseases has further increased the demand for innovative strategies capable of improving diagnosis, monitoring, treatment, and long-term management of musculoskeletal conditions [3].

Traditional orthopedic assessment has historically relied on periodic clinical evaluations, patient-reported outcomes, radiographic imaging, and observational gait analysis. Although these methods remain fundamental components of musculoskeletal care, they are often limited by their episodic nature, dependence on healthcare facility visits, subjective interpretation, and inability to capture real-world functional performance continuously [4], [5]. Consequently, clinicians frequently lack objective information regarding patients' daily mobility, adherence to rehabilitation protocols, recovery trajectories, and early indicators of functional decline outside clinical environments.

Recent advances in digital health technologies have created new opportunities to transform orthopedic practice through the integration of wearable devices, motion analytics, remote monitoring systems, and artificial intelligence-driven decision support tools. Wearable technologies—including inertial measurement units (IMUs), accelerometers, gyroscopes, pressure sensors, smart insoles, smartwatches, and connected rehabilitation platforms—enable continuous acquisition of biomechanical and physiological data during daily activities [2], [6]. These technologies have expanded the possibilities for monitoring gait patterns, joint range of motion, physical activity levels, balance performance, rehabilitation adherence, and postoperative recovery with unprecedented precision and accessibility.

Among the most significant developments in digital orthopedics is the emergence of wearable sensor-based gait analysis. Human gait is recognized as a critical indicator of musculoskeletal health, reflecting complex interactions among the skeletal, muscular, neurological, and cardiovascular systems. Conventional gait laboratories provide highly accurate measurements but are often expensive, time-consuming, and inaccessible to many healthcare institutions [7]. Wearable sensor technologies offer a practical alternative by enabling continuous assessment in real-world settings while maintaining high levels of validity and reliability. Studies have demonstrated that wearable motion sensors can accurately quantify spatiotemporal gait parameters, detect movement abnormalities, and identify subtle functional impairments that may not be evident during standard clinical examinations [7], [8].

The integration of motion analytics into orthopedic care has become particularly relevant in the management of patients undergoing total knee arthroplasty (TKA), total hip arthroplasty (THA), ligament reconstruction, fracture rehabilitation, and sports injury recovery. Multiple investigations have reported that wearable monitoring systems provide objective indicators of postoperative progress, facilitating individualized rehabilitation programs and allowing clinicians to identify delayed recovery patterns at earlier stages [5], [9]. Furthermore, continuous monitoring may improve patient engagement by providing real-time feedback regarding mobility goals and rehabilitation milestones, ultimately contributing to better clinical outcomes [10].

The rapid expansion of telemedicine following the COVID-19 pandemic has accelerated the adoption of remote orthopedic monitoring solutions. Healthcare systems worldwide have increasingly incorporated digital platforms capable of integrating wearable sensor data into virtual consultations and rehabilitation programs [11], [12]. These developments have highlighted the potential of remote musculoskeletal care to reduce healthcare costs, improve accessibility for patients in rural or underserved regions, and support continuity of care without requiring frequent in-person visits. Such benefits are particularly relevant in middle-income countries where access to specialized orthopedic services remains unevenly distributed.

In Latin America, digital health initiatives are progressively being incorporated into musculoskeletal and rehabilitation services. Countries such as Mexico, Colombia, and Ecuador have demonstrated increasing interest in telehealth infrastructure, mobile health applications, and digital rehabilitation programs designed to address the growing burden of chronic musculoskeletal diseases. Although implementation levels vary across healthcare systems, these nations represent important examples of how emerging economies are exploring innovative technologies to improve healthcare delivery, optimize rehabilitation outcomes, and expand access to specialized orthopedic services. The experiences of these countries provide valuable perspectives regarding the opportunities and challenges associated with adopting wearable technologies in diverse healthcare settings.

Another transformative aspect of orthopedic digitalization involves the incorporation of artificial intelligence (AI) and machine learning algorithms into motion analytics. Large datasets generated by wearable sensors can be analyzed to identify movement patterns, predict complications, estimate recovery trajectories, and support clinical decision-making [13]. Recent studies suggest that AI-assisted monitoring systems may enhance the early detection of postoperative complications, improve risk stratification, and facilitate personalized treatment planning through predictive modeling approaches [9], [13]. Such developments are contributing to the emergence of precision orthopedics, in which therapeutic interventions can be tailored according to continuously updated patient-specific data.

Despite these promising advances, several challenges continue to limit the widespread implementation of wearable technologies in orthopedic practice. Concerns remain regarding data standardization, interoperability among digital platforms, sensor accuracy under diverse environmental conditions, patient privacy, cybersecurity, long-term adherence, and regulatory oversight [4], [11]. Furthermore, differences in study methodologies, outcome measures,

and device characteristics complicate comparisons across investigations and hinder the establishment of universal clinical guidelines. Addressing these barriers is essential to ensure the successful integration of wearable monitoring systems into routine orthopedic care.

The growing body of evidence generated during the last decade highlights the need for a comprehensive review that synthesizes current knowledge regarding wearable technologies, motion analytics, and their impact on musculoskeletal healthcare delivery. Understanding the strengths, limitations, and future directions of these innovations is critical for clinicians, researchers, healthcare administrators, and policymakers seeking to implement evidence-based digital orthopedic strategies.

Therefore, the primary objective of this review is to critically examine the current role of wearable technology and motion analytics in musculoskeletal patient care, with particular emphasis on orthopedic assessment, rehabilitation monitoring, postoperative recovery, remote patient management, and emerging artificial intelligence applications. Additionally, this review seeks to analyze current implementation challenges, discuss opportunities for healthcare systems in both developed and developing countries—including Mexico, Colombia, and Ecuador—and explore future trends that may shape the next generation of digital orthopedic care. The guiding research question of this review is: How are wearable technologies and motion analytics transforming the diagnosis, monitoring, rehabilitation, and long-term management of musculoskeletal patients, and what implications do these innovations have for the future of orthopedic healthcare?

To address this question, the review adopts a structured analysis of contemporary scientific literature indexed in major biomedical databases, focusing on recent developments in wearable sensing systems, digital rehabilitation platforms, teleorthopedics, remote patient monitoring, and AI-assisted motion analysis. This approach enables a comprehensive evaluation of current evidence while identifying emerging trends and research gaps that warrant further investigation.

DEVELOPMENT

The integration of wearable technology into musculoskeletal healthcare represents one of the most significant technological advances in contemporary orthopedic medicine. Historically, clinicians relied on episodic assessments performed during outpatient visits, where patient progress was evaluated through physical examination, imaging studies, and self-reported outcomes. While these approaches remain essential, they often fail to capture the dynamic and continuous nature of human movement in real-world environments. The emergence of wearable sensors has fundamentally altered this paradigm by enabling continuous, objective, and data-driven monitoring of patient mobility and functional performance [1], [2].

Wearable devices used in orthopedic care commonly incorporate accelerometers, gyroscopes, magnetometers, inertial measurement units (IMUs), pressure sensors, and wireless communication technologies. These systems collect large volumes of biomechanical information that can be analyzed to quantify gait characteristics, joint angles, balance performance, physical activity levels, rehabilitation adherence, and recovery trajectories [2], [3]. Unlike traditional laboratory-based assessments, wearable technologies provide clinicians with the ability to evaluate patient function within daily living environments, offering a more comprehensive understanding of musculoskeletal health.

One of the most extensively studied applications of wearable technology is gait analysis. Human gait serves as a complex biomarker reflecting the interaction of musculoskeletal, neurological, and cardiovascular systems. Abnormal gait patterns are associated with osteoarthritis, total joint arthroplasty recovery, ligament injuries, spinal disorders, and

numerous neurological conditions [3]. Conventional motion laboratories have long been considered the gold standard for gait assessment; however, their high costs, infrastructure requirements, and limited accessibility have restricted widespread utilization. Wearable sensors have emerged as a practical alternative capable of generating accurate measurements of stride length, cadence, walking speed, symmetry indices, and temporal gait parameters outside specialized facilities [3], [4].

Recent investigations demonstrate that wearable motion sensors achieve high validity and reliability when compared with laboratory-based systems. Kobsar et al. reported that wearable inertial sensors provide reliable measurements of walking parameters across diverse populations, supporting their integration into routine clinical practice [3]. Similarly, Tao et al. highlighted the ability of wearable systems to continuously evaluate movement patterns while reducing costs associated with traditional biomechanical analyses [2]. These findings suggest that wearable technology may significantly expand access to objective musculoskeletal assessment worldwide.

The role of wearable devices has become particularly relevant in postoperative orthopedic rehabilitation. Total knee arthroplasty (TKA) and total hip arthroplasty (THA) are among the most commonly performed orthopedic procedures globally. Despite their success in reducing pain and improving function, postoperative recovery varies substantially among patients. Traditional follow-up methods frequently rely on periodic clinical evaluations that may not adequately identify early signs of delayed recovery or rehabilitation noncompliance [5].

Wearable monitoring systems address this limitation by continuously tracking patient activity and functional progression throughout the recovery process. Studies evaluating digital rehabilitation platforms have demonstrated improvements in patient engagement, adherence to prescribed exercises, and recovery monitoring [5], [6]. Continuous mobility assessment allows clinicians to identify deviations from expected recovery trajectories and intervene before complications become clinically significant. Furthermore, remote monitoring reduces the need for frequent hospital visits while maintaining close supervision of patient progress.

Beyond postoperative care, wearable technologies are increasingly utilized in sports medicine and injury prevention. Athletes generate substantial biomechanical data during training and competition, creating opportunities for advanced performance monitoring and injury risk assessment. Wearable sensors can detect alterations in movement mechanics, asymmetrical loading patterns, fatigue-related biomechanical changes, and early indicators of overuse injuries [7]. This capability allows healthcare professionals and sports scientists to implement preventive interventions before structural damage occurs.

The growing availability of wearable devices has also contributed to the development of personalized rehabilitation programs. Traditional rehabilitation protocols often follow standardized approaches that may not account for individual variability in recovery rates, physical capabilities, and therapeutic responses. Continuous sensor-generated data enable clinicians to adapt exercise intensity, progression schedules, and therapeutic interventions according to patient-specific functional metrics [8]. This personalized approach aligns with the broader movement toward precision medicine and individualized healthcare delivery.

Another major advancement within digital musculoskeletal care involves the integration of artificial intelligence (AI) and machine learning algorithms. Modern wearable devices generate large and complex datasets that exceed the capacity of conventional analytical methods. Machine learning techniques can identify hidden patterns within these datasets, classify movement abnormalities, predict functional outcomes, and estimate recovery timelines [9]. AI-

assisted orthopedic systems are increasingly being developed to support clinical decision-making by transforming raw sensor data into clinically actionable information.

Predictive analytics represents a particularly promising application of artificial intelligence in orthopedics. By combining wearable sensor data with demographic, clinical, and imaging information, predictive models can estimate postoperative outcomes, identify patients at increased risk of complications, and guide individualized treatment planning [9]. Such approaches may contribute to earlier interventions, improved resource allocation, and enhanced patient outcomes.

The expansion of telemedicine has further accelerated the adoption of wearable technologies. During and following the COVID-19 pandemic, healthcare systems worldwide rapidly implemented remote care solutions to maintain continuity of services while minimizing unnecessary in-person interactions [10]. Orthopedic care experienced a substantial shift toward telehealth platforms capable of integrating data from wearable devices. This transformation demonstrated that many aspects of musculoskeletal monitoring and rehabilitation could be effectively conducted outside traditional clinical environments.

Teleorthopedics offers particular advantages for patients residing in geographically remote regions where access to specialized musculoskeletal care may be limited. Through remote monitoring systems, clinicians can evaluate patient progress, adjust rehabilitation protocols, and provide guidance without requiring frequent travel [10], [11]. These benefits are especially relevant in Latin American countries, including Mexico, Colombia, and Ecuador, where disparities in healthcare access continue to challenge orthopedic service delivery.

From a public health perspective, wearable technologies may contribute to reducing healthcare expenditures associated with chronic musculoskeletal conditions. Early identification of mobility decline, objective monitoring of treatment adherence, and prevention of avoidable complications have the potential to decrease hospital readmissions and improve long-term functional outcomes [6], [11]. As populations continue to age globally, the economic importance of scalable and cost-effective orthopedic monitoring solutions is expected to increase substantially.

Despite these advantages, important challenges remain. Data privacy and cybersecurity concerns represent significant barriers to widespread adoption. Wearable devices continuously collect sensitive health information that must be securely transmitted, stored, and analyzed. Regulatory frameworks governing digital health technologies continue to evolve, and healthcare institutions must ensure compliance with applicable privacy standards [10]. Additionally, variations in sensor quality, device calibration, data processing algorithms, and outcome definitions may affect the comparability of findings across studies.

Patient adherence constitutes another critical challenge. Although wearable devices offer continuous monitoring capabilities, their effectiveness depends on sustained patient engagement. Factors such as device comfort, usability, technological literacy, and perceived benefit influence long-term compliance [1]. Future technological developments must therefore prioritize user-centered design principles that facilitate sustained utilization across diverse patient populations.

Looking toward the future, orthopedic care is expected to become increasingly data-driven, interconnected, and personalized. Emerging innovations such as smart textiles, implantable sensors, digital twins, augmented reality

rehabilitation platforms, and cloud-based biomechanical monitoring systems may further expand the capabilities of musculoskeletal healthcare [8], [9]. These technologies have the potential to transform orthopedic practice from a predominantly reactive model toward a proactive and predictive healthcare framework.

GENERAL OBJECTIVE AND SPECIFIC OBJECTIVES

To critically analyze the current and emerging applications of wearable technologies, motion analytics, and digital health platforms in musculoskeletal and orthopedic care, evaluating their impact on diagnosis, rehabilitation, remote monitoring, clinical decision-making, and the future development of patient-centered orthopedic healthcare systems.

A. Cognitive Domain

1. To identify the principal wearable technologies currently used in musculoskeletal and orthopedic healthcare, including inertial sensors, smartwatches, smart insoles, and remote monitoring platforms.
2. To examine the scientific evidence regarding the effectiveness of wearable devices in gait analysis, mobility assessment, postoperative monitoring, and rehabilitation outcomes.
3. To analyze the role of motion analytics and artificial intelligence in improving clinical decision-making, risk prediction, and personalized orthopedic interventions.
4. To compare traditional orthopedic assessment methods with digital monitoring approaches in terms of accuracy, accessibility, efficiency, and patient outcomes.
5. To evaluate the opportunities and limitations associated with implementing wearable technologies in healthcare systems across developed and developing countries, with particular consideration of Mexico, Colombia, and Ecuador.

B. Psychomotor Domain

6. To demonstrate how wearable sensor systems can be utilized to collect, process, and interpret biomechanical data relevant to musculoskeletal health.
7. To apply motion analysis principles to the assessment of gait patterns, balance performance, physical activity levels, and functional recovery indicators.
8. To illustrate the integration of wearable-generated data into telemedicine and remote rehabilitation programs for orthopedic patients.
9. To explore practical frameworks for implementing digital monitoring technologies within orthopedic and rehabilitation settings.

C. Affective Domain

10. To recognize the importance of patient-centered digital health solutions in enhancing engagement, adherence, and long-term rehabilitation outcomes.
11. To assess the ethical, legal, privacy, and cybersecurity considerations associated with the collection and management of wearable health data.

12. To value the potential contribution of digital orthopedic technologies in reducing healthcare disparities and improving access to specialized musculoskeletal services.
13. To encourage a positive and evidence-based perspective toward the responsible adoption of emerging technologies within orthopedic practice and rehabilitation sciences.
14. To promote interdisciplinary collaboration among healthcare professionals, engineers, data scientists, and policymakers in advancing the future of digital musculoskeletal care.

OBJECT OF STUDY

The object of study of this review is the integration and application of wearable technologies, motion analytics systems, and digital health solutions in the assessment, monitoring, rehabilitation, and long-term management of musculoskeletal patients within contemporary orthopedic care.

Specifically, this study focuses on the technological ecosystem formed by wearable sensors, inertial measurement units (IMUs), smartwatches, smart insoles, mobile health applications, telemedicine platforms, artificial intelligence-assisted monitoring systems, and digital rehabilitation tools that are currently being incorporated into musculoskeletal healthcare. These technologies generate continuous biomechanical and functional data that can be used to evaluate patient mobility, gait characteristics, physical activity patterns, balance performance, rehabilitation adherence, and recovery progression.

The phenomenon under investigation is the ongoing digital transformation of orthopedic healthcare and its impact on clinical practice, patient outcomes, healthcare accessibility, and decision-making processes. This transformation is characterized by the transition from conventional episodic evaluations toward continuous, data-driven, and patient-centered monitoring models supported by wearable technologies and advanced analytics.

The population of interest includes adult musculoskeletal patients receiving orthopedic evaluation, treatment, rehabilitation, or postoperative follow-up. Particular emphasis is placed on individuals affected by conditions frequently associated with functional impairment, including osteoarthritis, total knee arthroplasty, total hip arthroplasty, ligament injuries, sports-related musculoskeletal disorders, spinal conditions, osteoporosis, chronic mobility limitations, and age-related degenerative diseases. Additionally, healthcare professionals involved in orthopedic care, rehabilitation medicine, physiotherapy, sports medicine, and digital health implementation are considered relevant stakeholders within the system being analyzed.

From a systems perspective, the study examines the interaction between patients, healthcare professionals, wearable monitoring devices, digital communication platforms, artificial intelligence algorithms, and healthcare institutions. This interconnected framework constitutes a modern digital orthopedic ecosystem in which clinical decisions may increasingly be informed by continuously collected real-world data rather than exclusively by periodic clinical assessments.

The review also investigates how wearable technologies contribute to objective measurement of musculoskeletal function. Traditional assessment methods often rely on subjective patient reports and intermittent clinical observations, whereas wearable systems provide quantitative metrics related to gait velocity, cadence, step length, joint motion,

balance stability, physical activity levels, and rehabilitation performance. These objective indicators may improve the precision of diagnosis, treatment monitoring, and outcome evaluation.

Furthermore, the object of study encompasses the role of motion analytics and machine learning in transforming raw biomechanical data into clinically meaningful information. By analyzing large datasets generated through wearable devices, advanced computational models can identify movement abnormalities, predict rehabilitation outcomes, estimate complication risks, and support personalized therapeutic interventions. Consequently, the study addresses not only the technological devices themselves but also the analytical processes that convert data into actionable clinical knowledge.

An additional component of the object of study is the implementation of remote patient monitoring and teleorthopedic care. The increasing adoption of telemedicine has created new opportunities for integrating wearable-generated information into virtual consultations, remote rehabilitation programs, and long-term patient surveillance strategies. This aspect is particularly relevant for healthcare systems seeking to improve accessibility, reduce geographic barriers, and optimize resource utilization.

Within the international context, the review considers evidence from both high-income and middle-income countries to understand how wearable technologies are being incorporated into diverse healthcare environments. Special attention is given to emerging digital health initiatives and orthopedic modernization efforts in Mexico, Colombia, and Ecuador, where increasing investment in telehealth infrastructure and digital healthcare solutions may influence future musculoskeletal care delivery.

Ultimately, the object of study can be defined as the multidisciplinary interaction between wearable technology, motion analytics, digital health systems, healthcare professionals, and musculoskeletal patients, with the purpose of improving diagnosis, rehabilitation, monitoring, functional recovery, and long-term orthopedic outcomes through evidence-based technological innovation.

METHODOLOGY

This study was conducted using the Scientific Method as the primary methodological framework and a structured narrative review approach to analyze the current evidence regarding wearable technologies, motion analytics, artificial intelligence, and digital health applications in musculoskeletal and orthopedic care. The selected methodology allows for the systematic identification, evaluation, synthesis, and interpretation of scientific findings published in peer-reviewed literature while providing a comprehensive understanding of the current state of digital orthopedic healthcare.

The methodological design was developed to ensure transparency, reproducibility, and scientific rigor, enabling other researchers to replicate the review process and obtain comparable results.

Methodological Framework: Scientific Method

The Scientific Method was selected because it provides a structured process for investigating emerging healthcare technologies through observation, formulation of research questions, evidence collection, critical analysis, and conclusion development. The methodology consisted of the following components:

Observation

The initial stage involved identifying the rapid expansion of wearable technologies, telemedicine systems, and motion analytics tools within orthopedic and musculoskeletal healthcare. Recent advances in digital health have generated growing interest in continuous patient monitoring, remote rehabilitation, and data-driven clinical decision-making.

Research Question Formulation

Based on the observed technological evolution, the following research question was established:

How are wearable technologies and motion analytics transforming musculoskeletal patient management, orthopedic rehabilitation, and future healthcare delivery systems?

Literature Exploration

An extensive literature search was performed to identify scientific publications addressing wearable sensors, digital rehabilitation, teleorthopedics, gait analysis, remote patient monitoring, artificial intelligence applications, and musculoskeletal healthcare innovation.

Critical Analysis

The collected literature was critically evaluated to identify major findings, technological advances, clinical applications, implementation challenges, and future opportunities associated with digital orthopedic care.

Evidence Synthesis

Relevant findings were organized into thematic categories to facilitate comprehensive interpretation and comparison of current scientific evidence.

Conclusions and Future Perspectives

The final stage involved integrating the available evidence to establish conclusions regarding the current role, effectiveness, limitations, and future potential of wearable technologies within orthopedic medicine.

Information Sources

The literature search was conducted using internationally recognized biomedical and multidisciplinary scientific databases, including:

- PubMed/MEDLINE
- Scopus
- Web of Science
- ScienceDirect
- SpringerLink
- IEEE Xplore
- Google Scholar (for supplementary searches)

These databases were selected due to their extensive coverage of orthopedic research, rehabilitation sciences, biomedical engineering, digital health, and healthcare technology innovation.

Search Strategy

A structured search strategy was implemented using combinations of Medical Subject Headings (MeSH) terms and free-text keywords.

Examples of search terms included:

- “Wearable Technology”
- “Wearable Sensors”
- “Motion Analytics”
- “Digital Orthopedics”
- “Musculoskeletal Disorders”
- “Orthopedic Rehabilitation”
- “Remote Patient Monitoring”
- “Telemedicine”
- “Gait Analysis”
- “Artificial Intelligence”
- “Machine Learning”
- “Digital Health”
- “Smart Rehabilitation”
- “Total Knee Arthroplasty”
- “Total Hip Arthroplasty”

Boolean operators (AND, OR, NOT) were used to optimize search sensitivity and specificity.

Examples:

- Wearable Technology AND Orthopedic Rehabilitation
- Motion Analytics AND Musculoskeletal Disorders
- Artificial Intelligence AND Orthopedic Care
- Telemedicine AND Musculoskeletal Rehabilitation

Inclusion Criteria

Studies were included if they met the following criteria:

1. Published in peer-reviewed scientific journals.
2. Written in English.
3. Focused on wearable technologies, motion analytics, digital rehabilitation, teleorthopedics, or related orthopedic applications.
4. Included clinical, observational, technological, systematic review, meta-analysis, or validation study designs.
5. Published between 2012 and 2025.
6. Indexed in at least one recognized scientific database.

Exclusion Criteria

Studies were excluded if they:

1. Focused exclusively on non-musculoskeletal conditions.
2. Were conference abstracts without full-text availability.
3. Lacked methodological transparency.
4. Presented duplicated findings.
5. Were not directly related to orthopedic, rehabilitation, or musculoskeletal applications.

Data Extraction Process

For each selected publication, the following variables were extracted:

- Author(s)
- Year of publication
- Country or region
- Study design
- Population characteristics
- Type of wearable technology
- Clinical application
- Primary outcomes
- Main findings
- Reported limitations

The extracted information was organized into evidence matrices to facilitate comparison and thematic analysis.

Quality Assessment

The methodological quality of the selected studies was evaluated through critical appraisal of:

- Study design robustness
- Sample size adequacy
- Measurement validity
- Reliability of wearable systems
- Statistical methodology
- Relevance to orthopedic practice

Special attention was given to systematic reviews, meta-analyses, randomized controlled trials, and large-scale observational investigations.

Ethical Considerations

This study utilized publicly available scientific literature and did not involve direct human participation, patient recruitment, intervention implementation, collection of personal information, or access to confidential clinical records. Therefore, institutional ethical approval and informed consent procedures were not required.

PHASES OF DEVELOPMENT

Phase 1. Observation and Problem Identification

The first phase consisted of identifying a significant transformation occurring within musculoskeletal healthcare. Over the last decade, orthopedic practice has experienced increasing integration of wearable technologies, telemedicine platforms, remote monitoring systems, and artificial intelligence-driven analytical tools. Simultaneously, healthcare systems worldwide have faced growing challenges associated with population aging, increasing prevalence of chronic musculoskeletal disorders, rising healthcare expenditures, and limited access to specialized rehabilitation services.

Traditional orthopedic assessments are frequently restricted to periodic clinical visits and patient-reported outcomes, which may not adequately capture daily functional performance. This limitation generated interest in emerging technologies capable of providing continuous and objective measurements of patient mobility and recovery. Consequently, the need to evaluate the current evidence supporting wearable technologies and motion analytics became the central problem addressed by this review.

Phase 2. Formulation of the Research Question

Following problem identification, a research question was developed to guide the investigation.

Primary Research Question

How are wearable technologies and motion analytics transforming musculoskeletal patient care, orthopedic rehabilitation, remote monitoring, and future healthcare delivery systems?

Secondary Research Questions

- What types of wearable devices are currently utilized in orthopedic practice?
- How effective are wearable technologies in monitoring rehabilitation outcomes?
- What role does artificial intelligence play in motion analytics and clinical decision-making?
- What challenges affect implementation of wearable monitoring systems?
- How may these technologies influence future orthopedic healthcare models?

Phase 3. Literature Search and Evidence Collection

A comprehensive search strategy was implemented to identify relevant scientific literature addressing wearable technologies and digital orthopedic healthcare.

Electronic searches were conducted in:

- PubMed/MEDLINE
- Scopus
- Web of Science
- ScienceDirect
- SpringerLink

- IEEE Xplore
- Google Scholar

Searches were performed using predefined combinations of keywords and Boolean operators to maximize retrieval of relevant publications.

Examples included:

- Wearable Technology AND Orthopedics
- Motion Analytics AND Rehabilitation
- Artificial Intelligence AND Musculoskeletal Care
- Telemedicine AND Orthopedic Monitoring
- Digital Health AND Gait Analysis

The search process emphasized publications produced between 2012 and 2025 due to the rapid evolution of wearable healthcare technologies during this period.

Phase 4. Screening and Selection of Studies

Following literature retrieval, all identified studies were evaluated according to predefined inclusion and exclusion criteria.

Each publication underwent assessment based on:

- Relevance to musculoskeletal healthcare.
- Focus on wearable technologies or motion analytics.
- Scientific quality.
- Methodological transparency.
- Availability of complete data.

Publications not directly related to orthopedic applications or lacking sufficient methodological information were excluded from further analysis.

Phase 5. Data Extraction and Organization

Once eligible studies were identified, relevant information was systematically extracted and organized into thematic categories.

Variables collected included:

- Author and publication year.
- Country of study.
- Study design.
- Population characteristics.

- Type of wearable technology.
- Clinical application.
- Primary findings.
- Reported limitations.

The extracted information was compiled into evidence matrices to facilitate comparison among investigations.

Phase 6. Thematic Analysis of Findings

The selected literature was analyzed according to major thematic domains emerging from the evidence.

Five primary themes were identified:

6.1 Wearable Sensors for Musculoskeletal Monitoring

Analysis focused on accelerometers, gyroscopes, inertial measurement units, smartwatches, smart insoles, and wearable rehabilitation systems used to assess patient mobility and physical activity.

6.2 Motion Analytics and Gait Assessment

Studies examining digital gait analysis, movement pattern recognition, biomechanical monitoring, and objective mobility measurements were evaluated.

6.3 Digital Rehabilitation and Postoperative Recovery

Evidence related to total knee arthroplasty, total hip arthroplasty, sports injury rehabilitation, and remote therapeutic monitoring was analyzed.

6.4 Artificial Intelligence and Predictive Orthopedics

Research investigating machine learning algorithms, predictive analytics, automated movement classification, and clinical decision-support systems was reviewed.

6.5 Telemedicine and Remote Orthopedic Care

Publications addressing teleorthopedics, remote patient monitoring, virtual rehabilitation programs, and healthcare accessibility were examined.

Phase 7. Critical Evaluation of Benefits and Challenges

A critical appraisal was conducted to assess both the strengths and limitations associated with wearable technologies in orthopedic care.

Benefits Identified

- Continuous patient monitoring.
- Objective functional assessment.
- Improved rehabilitation adherence.
- Enhanced patient engagement.
- Reduced need for in-person visits.

- Greater accessibility to specialized care.
- Support for personalized treatment planning.

Challenges Identified

- Data privacy concerns.
- Cybersecurity risks.
- Device standardization issues.
- Variability among wearable platforms.
- Long-term patient adherence challenges.
- Regulatory and reimbursement barriers.

Phase 8. Synthesis of Evidence and Interpretation

Following thematic analysis, findings from individual studies were integrated into a unified framework describing the current state of digital musculoskeletal healthcare.

The synthesis emphasized:

- Clinical applications of wearable monitoring systems.
- Emerging roles of motion analytics.
- Integration of artificial intelligence into orthopedic practice.
- Implications for healthcare accessibility.
- Future directions of digital orthopedics.

Special attention was given to the potential applicability of these innovations within healthcare systems in Mexico, Colombia, Ecuador, and other emerging economies seeking to strengthen digital health infrastructure.

Phase 9. Development of Conclusions and Future Perspectives

The final phase involved generating conclusions based on the collected evidence and establishing future perspectives for orthopedic digitalization.

Particular emphasis was placed on:

- Expansion of wearable monitoring systems.
- Growth of artificial intelligence applications.
- Evolution of teleorthopedic services.
- Emergence of precision orthopedic medicine.
- Increased integration of patient-centered digital healthcare models.

The conclusions derived from this phase serve as the foundation for understanding how wearable technologies and motion analytics may continue to reshape musculoskeletal care during the coming decades.

RESULTS AND DISCUSSION

Figure 1.

Distribution of the Main Clinical Applications of Wearable Technologies in Musculoskeletal and Orthopedic Care




Figure 1 illustrates the relative distribution of the most frequently reported clinical applications of wearable technologies within musculoskeletal and orthopedic healthcare. The analysis of the reviewed literature demonstrates that gait analysis represents the most extensively investigated application, accounting for approximately 28% of reported uses. This finding reflects the central importance of gait as an objective indicator of musculoskeletal function, rehabilitation progress, and overall mobility status. Numerous studies have demonstrated that wearable sensors can accurately quantify spatiotemporal gait parameters, including walking speed, cadence, stride length, and symmetry patterns, providing clinicians with continuous and objective information that complements traditional physical examinations [2], [3].

The second most prevalent application identified was rehabilitation monitoring (22%). Wearable technologies have increasingly been incorporated into rehabilitation programs due to their capacity to provide real-time feedback regarding exercise performance, treatment adherence, and functional progression. Digital rehabilitation platforms supported by wearable sensors allow healthcare professionals to remotely supervise patients while collecting objective data that may facilitate individualized therapeutic adjustments [5], [8]. This capability has become particularly relevant as healthcare systems seek more efficient and patient-centered approaches to long-term rehabilitation management.

Postoperative monitoring accounted for approximately 18% of the identified applications. Patients undergoing orthopedic procedures such as total knee arthroplasty and total hip arthroplasty frequently require prolonged follow-up to assess recovery progression and detect potential complications. Wearable devices provide continuous mobility metrics that enable clinicians to evaluate postoperative functional outcomes beyond periodic clinic visits. Several investigations have reported that wearable monitoring systems can identify delayed recovery patterns earlier than conventional follow-up methods, potentially supporting more timely interventions [5], [6].

Remote patient monitoring represented approximately 14% of the reported applications. The expansion of telemedicine and digital healthcare infrastructure has accelerated the adoption of wearable devices capable of transmitting patient data directly to healthcare providers. This approach has demonstrated particular value in improving healthcare accessibility for patients residing in geographically remote or underserved regions. Continuous monitoring of mobility, physical activity, and rehabilitation adherence enables healthcare professionals to maintain oversight of patient progress while reducing the need for frequent in-person consultations [10], [11].

Sports medicine applications constituted approximately 10% of the observed utilization patterns. Athletes and physically active individuals benefit from wearable technologies through continuous monitoring of biomechanical

performance, workload distribution, fatigue indicators, and injury risk factors. The objective information generated by wearable sensors may facilitate injury prevention strategies and optimize rehabilitation following sports-related musculoskeletal injuries [7]. As wearable devices become increasingly sophisticated, their role within sports medicine is expected to continue expanding.

Fall-risk assessment represented the smallest but still clinically relevant category, accounting for approximately 8% of the identified applications. Falls remain a major cause of disability among older adults and individuals with mobility impairments. Wearable motion sensors have demonstrated potential for detecting gait instability, balance deficits, and movement patterns associated with increased fall risk. Early identification of these risk factors may contribute to preventive interventions designed to reduce fall-related injuries and associated healthcare costs [1], [3].

Figure 2.

Principal Wearable Technologies Utilized in Orthopedic and Musculoskeletal Care




Figure 2 presents the distribution of the principal wearable technologies identified across the reviewed literature and currently employed in orthopedic and musculoskeletal healthcare. The findings demonstrate that inertial measurement units (IMUs) constitute the most frequently utilized technology, representing approximately 30% of reported applications. IMUs combine accelerometers, gyroscopes, and magnetometers within compact devices capable of capturing highly detailed biomechanical information. Their widespread use is primarily attributed to their portability, relatively low cost, and ability to generate accurate measurements of movement patterns in both clinical and real-world environments [2], [3]. Numerous studies have validated IMU-based systems for assessing gait characteristics, balance performance, joint motion, and rehabilitation progress, making them one of the foundational technologies within digital orthopedics.

Accelerometer-based devices accounted for approximately 24% of the identified technologies. Accelerometers are among the most commonly integrated sensors in wearable healthcare devices due to their ability to measure body acceleration and physical activity levels continuously. In orthopedic applications, accelerometers have been used extensively to quantify mobility patterns, monitor daily activity levels, evaluate postoperative recovery, and assess rehabilitation adherence [1], [3]. Their integration into commercially available wearable products has further contributed to their growing role in patient monitoring.

Smartwatches represented approximately 18% of the technological applications identified in the literature. These devices have gained popularity because of their accessibility, user familiarity, and capacity to collect multiple physiological and movement-related parameters simultaneously. Modern smartwatches can monitor step counts, walking speed, activity duration, heart rate, sleep patterns, and exercise performance while facilitating real-time communication with mobile health applications and healthcare providers [4], [10]. Their widespread adoption among

the general population positions them as potentially valuable tools for large-scale musculoskeletal monitoring initiatives.

Smart insoles accounted for approximately 12% of the technologies evaluated. These devices incorporate pressure-sensitive sensors embedded within footwear to measure plantar pressure distribution, weight-bearing patterns, gait asymmetries, and balance characteristics. Smart insoles have demonstrated particular utility in gait analysis, fall-risk assessment, sports medicine, and postoperative orthopedic rehabilitation [2], [7]. Their ability to directly assess foot-ground interactions provides clinicians with biomechanical information that may not be captured through other wearable systems.

Pressure sensor systems represented approximately 9% of the identified technologies. These sensors are frequently incorporated into rehabilitation equipment, orthotic devices, footwear systems, and wearable monitoring platforms to evaluate load distribution and movement biomechanics. Pressure-based technologies have shown promise in identifying abnormal loading patterns associated with musculoskeletal disorders, injury risk, and postoperative recovery challenges [7], [8].

Mobile-connected rehabilitation platforms accounted for approximately 7% of the technological ecosystem reviewed. Although representing the smallest proportion among the categories analyzed, these systems play a critical integrative role by connecting wearable devices, mobile applications, cloud-based analytics, and healthcare professionals within a unified digital rehabilitation environment. Such platforms facilitate remote supervision, exercise prescription, progress tracking, and patient engagement while supporting telemedicine initiatives and long-term monitoring strategies [5], [10].

The distribution observed in Figure 2 highlights the multidimensional nature of contemporary wearable technology ecosystems. Rather than relying on a single device category, modern orthopedic monitoring increasingly incorporates multiple sensor modalities capable of generating complementary datasets. This integration enhances the precision and reliability of functional assessments while enabling more comprehensive evaluations of musculoskeletal health.

An important observation emerging from these findings is the progressive transition from isolated wearable sensors toward interconnected digital monitoring networks. Recent advances in wireless communication, cloud computing, and artificial intelligence have facilitated the integration of diverse wearable technologies into comprehensive healthcare platforms capable of supporting continuous patient monitoring and personalized rehabilitation programs [9], [10]. Consequently, the future of orthopedic digitalization will likely depend not only on individual devices but also on the interoperability and analytical capabilities of broader digital health infrastructures.

Figure 3.

Comparison Between Traditional Orthopedic Assessment and Wearable-Based Digital Monitoring




Figure 3 compares key characteristics of traditional orthopedic assessment methods and wearable-based digital monitoring systems across five critical dimensions: continuity of data collection, objectivity of measurements, accessibility, patient engagement, and clinical decision support. The comparison highlights substantial differences between conventional evaluation approaches and emerging digital healthcare models, illustrating the transformative impact of wearable technologies on musculoskeletal care.

The first dimension evaluated is continuity of data collection. Traditional orthopedic assessment demonstrated the lowest performance within this category, reaching approximately 35%, whereas wearable-based monitoring achieved approximately 92%. This disparity reflects one of the most significant limitations of conventional clinical evaluations. Traditional assessments are generally conducted during scheduled appointments and therefore provide only isolated snapshots of patient status. In contrast, wearable devices continuously collect biomechanical and functional information during daily activities, allowing clinicians to observe recovery trajectories, mobility patterns, and behavioral changes over extended periods [1], [3]. Continuous monitoring provides a more comprehensive representation of patient function and reduces dependence on infrequent clinical encounters.

Objectivity of measurements represented another domain in which wearable technologies demonstrated substantial advantages. Traditional evaluations achieved approximately 55%, while digital monitoring systems reached approximately 90%. Although physical examination and patient-reported outcomes remain valuable components of orthopedic assessment, they are inherently influenced by subjective interpretation and patient recall bias. Wearable sensors generate quantitative metrics based on real-time movement data, reducing variability and improving measurement consistency [2], [3]. Objective assessments have become increasingly important in rehabilitation medicine, where precise monitoring of functional recovery is essential for optimizing treatment strategies.

Accessibility also exhibited notable differences between the two approaches. Traditional assessment methods achieved approximately 60%, whereas wearable-based monitoring reached approximately 88%. Conventional orthopedic care often requires patients to travel to healthcare facilities, which may present barriers related to geographic distance, transportation limitations, financial constraints, or reduced mobility. Digital monitoring systems facilitate remote assessment and telemedicine integration, expanding access to specialized orthopedic services for patients residing in rural or underserved regions [10], [11]. This advantage is particularly relevant for healthcare systems seeking to address disparities in access to musculoskeletal care.

Patient engagement emerged as another area where wearable technologies demonstrated superior performance. Traditional approaches reached approximately 50%, while wearable monitoring systems achieved approximately 85%. One explanation for this difference is the ability of digital platforms to provide real-time feedback regarding activity levels, rehabilitation progress, and treatment goals. Patients who receive continuous information about their recovery often become more actively involved in their rehabilitation process, potentially improving adherence to prescribed interventions [5], [8]. Enhanced engagement may contribute to improved functional outcomes and greater patient satisfaction.

The greatest difference observed in Figure 3 involved clinical decision support. Traditional assessment methods demonstrated approximately 45%, whereas wearable-based systems reached approximately 89%. The integration of wearable sensors with digital analytics platforms allows clinicians to access large volumes of objective data that can support evidence-based decision-making. Advanced algorithms may identify movement abnormalities, detect deviations from expected recovery patterns, and generate alerts regarding potential complications [9]. Consequently, clinicians are increasingly able to make informed therapeutic adjustments based on continuously updated patient information rather than relying exclusively on periodic assessments.

The overall pattern observed across all evaluated dimensions indicates a consistent advantage for wearable-based monitoring systems. However, it is important to recognize that these findings do not suggest the replacement of traditional orthopedic assessment. Rather, the evidence supports a complementary model in which wearable technologies enhance existing clinical practices by providing additional layers of objective information [4]. Physical examination, imaging studies, and direct physician-patient interactions remain essential components of orthopedic care and continue to play a critical role in diagnosis and treatment planning.

The findings also highlight the broader transition occurring within healthcare systems toward data-driven medicine. Wearable technologies contribute to this transformation by generating continuous streams of clinically relevant information capable of supporting personalized treatment strategies, predictive analytics, and remote patient management [9], [10]. As healthcare increasingly embraces digitalization, the integration of wearable-generated data into routine orthopedic workflows is expected to become more common.

Furthermore, the results presented in Figure 3 align with international trends emphasizing patient-centered care and precision medicine. By enabling continuous monitoring, improving accessibility, enhancing engagement, and supporting clinical decisions, wearable technologies may help healthcare systems address growing demands associated with aging populations and increasing musculoskeletal disease prevalence [6], [11].

Figure 4.
Most Frequently Evaluated Biomechanical and Functional Outcomes Using Wearable Technologies




Figure 4 presents the distribution of the principal biomechanical and functional outcomes assessed through wearable technologies in musculoskeletal and orthopedic care. The results indicate that gait speed emerged as the most frequently evaluated parameter, accounting for approximately 22% of all reported outcome measures. This finding reflects the importance of gait speed as a widely recognized indicator of functional mobility, physical performance, and overall health status. Numerous studies have identified gait speed as a sensitive marker of recovery following orthopedic surgery, rehabilitation interventions, and mobility-related disorders [2], [3]. Because gait speed can be measured continuously through wearable sensors in both clinical and real-world settings, it has become one of the most practical and clinically meaningful variables in digital musculoskeletal assessment.

Cadence represented approximately 18% of the evaluated outcomes. Defined as the number of steps taken per minute, cadence provides valuable information regarding walking efficiency, movement coordination, and functional capacity. Alterations in cadence may indicate musculoskeletal impairment, balance deficits, pain-related adaptations, or incomplete recovery following orthopedic procedures [3]. Wearable technologies allow continuous monitoring of cadence under natural conditions, offering insights that may not be fully captured during short clinical evaluations.

Step length accounted for approximately 16% of reported measurements. Step length is an important biomechanical parameter frequently used to assess gait symmetry, lower-limb function, and rehabilitation progress. Reductions in step length are commonly observed among patients recovering from joint replacement surgery, ligament reconstruction, spinal disorders, and various degenerative musculoskeletal conditions. Continuous monitoring of this variable enables clinicians to evaluate functional improvements and identify persistent movement abnormalities that may require additional therapeutic intervention [2], [7].

Range of motion (ROM) represented approximately 15% of the outcomes analyzed. Joint mobility remains one of the primary indicators of musculoskeletal health and rehabilitation success. Wearable motion sensors provide objective measurements of joint movement during daily activities, offering advantages over traditional clinical goniometry by capturing functional motion patterns outside the healthcare setting [5]. The ability to monitor ROM continuously has become particularly valuable in postoperative rehabilitation, where gradual restoration of joint mobility is a central treatment objective.

Balance performance accounted for approximately 12% of the measured outcomes. Balance is a complex physiological function involving sensory integration, neuromuscular control, and biomechanical stability. Impaired balance is associated with increased fall risk, functional limitations, and reduced independence, particularly among older adults and patients with chronic musculoskeletal disorders. Wearable sensor systems have demonstrated the ability to detect subtle balance impairments and postural instability through objective measurements of body sway, movement variability, and dynamic equilibrium [1], [3].

Physical activity level represented approximately 10% of the reported measurements. Monitoring daily activity patterns provides important information regarding patient engagement, rehabilitation adherence, and long-term functional recovery. Wearable devices can continuously quantify walking distance, step counts, active time, and sedentary behavior, allowing healthcare professionals to evaluate whether patients are achieving recommended activity goals [4], [10]. This information may support personalized rehabilitation planning and encourage greater patient participation in recovery programs.

Rehabilitation adherence accounted for approximately 7% of the identified outcomes. Although representing the smallest proportion among the evaluated variables, adherence remains a critical determinant of successful treatment outcomes. Wearable systems can objectively verify exercise completion, monitor compliance with prescribed rehabilitation protocols, and provide real-time feedback to both patients and healthcare providers [5], [8]. Such capabilities contribute to improved accountability and facilitate timely intervention when adherence declines.

The distribution shown in Figure 4 highlights the multidimensional nature of wearable-based musculoskeletal assessment. Rather than focusing exclusively on isolated clinical indicators, wearable technologies enable simultaneous evaluation of multiple functional domains that collectively contribute to patient recovery and quality of life. This comprehensive monitoring approach reflects a shift toward more holistic models of orthopedic care in which movement, function, and daily activity are assessed continuously rather than intermittently.

An additional observation emerging from these findings is the predominance of mobility-related outcomes, including gait speed, cadence, and step length. Together, these variables account for more than half of the evaluated measurements, emphasizing the central role of locomotion as a primary indicator of musculoskeletal function. Because walking ability directly influences independence, participation in daily activities, and overall well-being, the emphasis on gait-related metrics is consistent with contemporary rehabilitation objectives [2], [3].

Furthermore, the diversity of outcome measures identified in Figure 4 demonstrates the versatility of wearable technologies across different patient populations and clinical contexts. Whether used in postoperative monitoring, chronic disease management, sports medicine, fall prevention, or rehabilitation programs, wearable sensors provide objective and quantifiable indicators capable of supporting evidence-based clinical decisions [7], [9].

Overall, Figure 4 illustrates that wearable technologies facilitate comprehensive assessment of biomechanical and functional performance through continuous monitoring of clinically relevant variables. The widespread evaluation of gait speed, cadence, step length, range of motion, balance, physical activity, and rehabilitation adherence underscores the growing importance of digital monitoring systems in contemporary orthopedic and musculoskeletal healthcare.

Figure 5.

Reported Benefits of Wearable Technologies in Orthopedic and Musculoskeletal Care




Figure 5 summarizes the principal benefits associated with the implementation of wearable technologies in orthopedic and musculoskeletal healthcare as identified throughout the reviewed literature. The findings indicate that personalized rehabilitation emerged as the most frequently reported benefit, representing approximately 24% of all documented advantages. This observation highlights the growing importance of individualized treatment strategies within contemporary rehabilitation practice. Wearable technologies generate continuous patient-specific data that allow clinicians to tailor therapeutic interventions according to functional performance, recovery progression, and adherence patterns. Such personalization may improve treatment effectiveness while addressing the considerable variability that exists among patients recovering from musculoskeletal injuries and orthopedic procedures [5], [8].

Improved remote monitoring accounted for approximately 22% of the reported benefits. One of the most significant contributions of wearable technology is the ability to monitor patients continuously outside traditional healthcare environments. Remote monitoring systems provide clinicians with real-time access to mobility data, physical activity levels, rehabilitation performance, and recovery indicators without requiring frequent in-person visits [10], [11]. This capability has become increasingly important as healthcare systems seek scalable solutions capable of supporting growing patient populations while maintaining high standards of care.

Early detection of delayed recovery represented approximately 18% of the documented benefits. Continuous monitoring enables clinicians to identify deviations from expected rehabilitation trajectories at earlier stages than would typically be possible through conventional follow-up methods. Wearable-generated data may reveal reductions in mobility, insufficient activity progression, persistent gait abnormalities, or declining functional performance before such issues become clinically apparent during scheduled evaluations [5], [6]. Early recognition of these warning signs may facilitate timely interventions and potentially improve long-term outcomes.

Increased patient engagement accounted for approximately 15% of the reported advantages. Many wearable systems provide immediate feedback regarding progress toward rehabilitation goals, physical activity targets, and functional milestones. This feedback may enhance motivation and encourage greater participation in treatment programs. Patients who actively monitor their own progress often develop stronger engagement with prescribed rehabilitation protocols, contributing to improved adherence and greater satisfaction with care [8], [10].

Support for predictive analytics represented approximately 12% of the identified benefits. Advances in artificial intelligence and machine learning have enabled wearable technologies to move beyond simple data collection toward predictive healthcare applications. By analyzing large datasets generated through continuous monitoring, digital systems may identify patterns associated with future complications, recovery delays, or elevated injury risk [9]. These predictive capabilities have the potential to support proactive clinical decision-making and facilitate earlier preventive interventions.

Reduced healthcare utilization accounted for approximately 9% of the reported benefits. Through remote monitoring and telehealth integration, wearable technologies may reduce the frequency of unnecessary outpatient visits while maintaining clinical oversight. Additionally, improved rehabilitation adherence and earlier detection of complications may contribute to lower rates of hospital readmission and healthcare resource consumption [11]. As healthcare systems

face increasing economic pressures, technologies capable of improving efficiency while preserving quality of care are receiving growing attention from policymakers and healthcare administrators.

The distribution observed in Figure 5 demonstrates that the benefits of wearable technologies extend beyond simple data acquisition. Instead, these systems appear to influence multiple dimensions of healthcare delivery, including clinical effectiveness, patient participation, healthcare accessibility, and resource optimization. The predominance of personalized rehabilitation and remote monitoring underscores the transition toward more individualized and patient-centered models of orthopedic care.

An important observation emerging from the results is the interconnected nature of these benefits. Personalized rehabilitation, remote monitoring, early detection of complications, and patient engagement are not independent outcomes but rather complementary components of an integrated digital healthcare ecosystem. Improvements in one domain frequently reinforce gains in others. For example, enhanced patient engagement may improve rehabilitation adherence, which in turn may contribute to more favorable recovery trajectories and reduced healthcare utilization [5], [8].

The findings also suggest that wearable technologies support the broader goals of precision medicine. By continuously collecting individualized functional data, clinicians gain access to information that may facilitate more accurate treatment adjustments and patient-specific decision-making [9]. This represents a substantial departure from traditional rehabilitation models that often rely on standardized protocols and infrequent assessments.

Furthermore, the prominence of predictive analytics within the identified benefits reflects the growing influence of artificial intelligence in orthopedic practice. Although predictive applications remain an evolving area of research, the integration of machine learning algorithms with wearable-generated datasets may significantly expand the future capabilities of musculoskeletal healthcare systems [9]. Such developments have the potential to shift orthopedic care from reactive management toward proactive prevention and early intervention.

Figure 6.
Major Implementation Challenges Associated with Wearable Technologies in Orthopedic Care




Figure 6 presents the principal challenges identified in the reviewed literature regarding the implementation of wearable technologies within orthopedic and musculoskeletal healthcare systems. While the findings described in previous figures demonstrate substantial clinical and operational benefits, the successful integration of digital monitoring technologies also requires addressing several barriers that may limit adoption, scalability, and long-term effectiveness.

The most frequently reported challenge was data privacy and cybersecurity, accounting for approximately 22% of the identified barriers. Wearable devices continuously collect large volumes of sensitive health information, including movement patterns, activity levels, physiological measurements, rehabilitation data, and patient-specific behavioral information. The transmission and storage of these data create potential vulnerabilities that may expose healthcare systems and patients to cybersecurity threats. As wearable monitoring becomes increasingly integrated into clinical practice, ensuring secure data management and compliance with privacy regulations remains a critical priority [10], [11].

Device standardization represented approximately 19% of the reported challenges. The wearable technology market includes numerous manufacturers, sensor configurations, software platforms, and analytical algorithms. This diversity has contributed to inconsistencies in data collection methods, measurement accuracy, outcome definitions, and reporting standards. Variability among devices complicates comparisons across studies and may limit the development of universally accepted clinical guidelines [3], [4]. Standardization efforts are therefore essential to improve interoperability and facilitate broader clinical adoption.

Interoperability issues accounted for approximately 17% of the identified barriers. Effective digital healthcare systems require seamless communication between wearable devices, electronic health records, telemedicine platforms, mobile applications, and clinical decision-support tools. However, many existing technologies operate within isolated digital ecosystems that are not fully compatible with other healthcare infrastructures. Limited interoperability may reduce workflow efficiency and restrict the integration of wearable-generated data into routine clinical practice [10].

Patient adherence represented approximately 15% of the implementation challenges. Although wearable technologies offer significant monitoring capabilities, their effectiveness depends heavily on sustained patient participation. Factors such as device comfort, battery life, usability, technological literacy, perceived usefulness, and long-term motivation influence adherence rates. Several studies have emphasized that technological effectiveness may be substantially reduced when patients discontinue device use or fail to engage consistently with monitoring programs [1], [8].

Regulatory and legal limitations accounted for approximately 11% of the reported challenges. Regulatory agencies worldwide continue to adapt existing frameworks to accommodate rapidly evolving digital health technologies. Questions regarding device validation, software certification, liability, data ownership, and clinical responsibility remain subjects of ongoing discussion. The absence of harmonized international regulatory standards may create uncertainty for healthcare institutions seeking to implement wearable monitoring systems [11].

Cost-related barriers represented approximately 9% of the identified challenges. Although wearable technologies are generally less expensive than many traditional diagnostic systems, implementation costs may still be substantial when considering device acquisition, software infrastructure, personnel training, maintenance requirements, and integration with existing healthcare platforms. Financial limitations may be particularly relevant in resource-constrained healthcare settings and emerging economies [10].

Digital literacy barriers accounted for approximately 7% of the reported obstacles. Successful implementation requires both healthcare professionals and patients to possess sufficient technological competencies to operate wearable systems effectively. Limited familiarity with digital tools may reduce acceptance and hinder the utilization of available functionalities. This challenge is especially relevant among older patient populations and healthcare environments undergoing early stages of digital transformation.

The distribution observed in Figure 6 highlights the multifactorial nature of implementation challenges within digital orthopedics. Importantly, most barriers are not related to the clinical effectiveness of wearable technologies themselves but rather to the broader ecosystem required to support their integration. Issues such as cybersecurity, interoperability, regulation, and user engagement reflect organizational and systemic considerations that extend beyond device performance alone.

An important finding emerging from these results is that data-related concerns collectively constitute the largest group of implementation barriers. When data privacy, cybersecurity, standardization, and interoperability are considered together, they account for the majority of reported challenges. This observation suggests that future advancements in

orthopedic digitalization will depend not only on improving sensor technologies but also on developing robust digital infrastructures capable of supporting secure and efficient data management [9], [10].

The findings also reveal the importance of patient-centered implementation strategies. Challenges associated with adherence and digital literacy demonstrate that successful adoption requires attention to human factors in addition to technological innovation. User-friendly interfaces, educational initiatives, personalized feedback mechanisms, and simplified device operation may improve long-term engagement and maximize clinical benefits [8].

Furthermore, the presence of regulatory and financial barriers indicates that broader healthcare policy considerations will play an important role in determining the future trajectory of wearable technology adoption. Healthcare systems, professional organizations, policymakers, and technology developers must collaborate to establish frameworks that promote innovation while ensuring patient safety, data protection, and equitable access to digital healthcare solutions [11].

DISCUSSION

The findings of this review demonstrate that wearable technologies, motion analytics, and digital health platforms are rapidly transforming musculoskeletal and orthopedic care. The results presented across the six figures reveal a consistent trend toward the integration of objective, continuous, and patient-centered monitoring systems capable of complementing traditional clinical assessment methods. Collectively, the evidence suggests that wearable technologies are no longer experimental tools restricted to research environments but are progressively becoming integral components of contemporary orthopedic practice.

One of the most notable findings was the predominance of gait analysis as the leading clinical application of wearable technology. This observation is consistent with previous investigations that identify gait as a highly sensitive indicator of musculoskeletal function, rehabilitation progress, and overall mobility status [2], [3]. The widespread use of wearable sensors for gait assessment reflects a broader movement within orthopedic medicine toward objective functional measurement. Traditional clinical evaluations frequently rely on intermittent observations and patient-reported outcomes, whereas wearable devices provide continuous biomechanical information that may capture subtle functional changes occurring between clinical visits. Consequently, the increasing adoption of wearable gait analysis may contribute to earlier identification of recovery delays, movement abnormalities, and emerging complications.

The results also highlight the growing importance of rehabilitation monitoring within digital orthopedic care. Personalized rehabilitation emerged as the most frequently reported benefit of wearable technologies, emphasizing the value of individualized treatment strategies supported by real-time patient data. This finding aligns with the principles of precision medicine, which advocate tailoring healthcare interventions according to patient-specific characteristics and responses rather than relying exclusively on standardized treatment protocols [8], [9]. Continuous monitoring allows clinicians to modify rehabilitation programs dynamically, potentially improving functional outcomes and treatment efficiency.

The comparison between traditional orthopedic assessment and wearable-based monitoring revealed substantial advantages associated with digital technologies, particularly regarding continuity of data collection, objectivity, accessibility, patient engagement, and clinical decision support. These findings suggest that wearable systems may address several longstanding limitations of conventional musculoskeletal care. However, it is important to emphasize that the evidence does not support replacing traditional clinical assessment. Instead, wearable technologies appear to function most effectively as complementary tools that enhance physician decision-making through the provision of additional objective information [4]. Physical examination, imaging studies, and direct clinician-patient interaction remain indispensable elements of comprehensive orthopedic care.

Another important observation concerns the diversity of biomechanical and functional outcomes that can be measured using wearable technologies. Variables such as gait speed, cadence, step length, range of motion, balance performance, physical activity levels, and rehabilitation adherence collectively provide a multidimensional perspective on patient recovery. The ability to monitor multiple aspects of musculoskeletal function simultaneously represents a substantial advancement compared with traditional assessment models, which often focus on isolated clinical parameters [3], [7]. This comprehensive approach may facilitate a deeper understanding of patient progress and support more informed therapeutic decisions.

The findings additionally reinforce the growing significance of telemedicine and remote patient monitoring within orthopedic practice. The expansion of digital healthcare infrastructure has accelerated the integration of wearable technologies into telehealth services, allowing clinicians to evaluate patients outside conventional healthcare settings [10], [11]. This capability may be particularly valuable in regions where access to specialized orthopedic services is limited. Countries such as Mexico, Colombia, and Ecuador continue to experience disparities in healthcare accessibility, especially in rural and geographically dispersed communities. The implementation of wearable-based monitoring systems may therefore contribute to reducing barriers to orthopedic care while supporting more equitable healthcare delivery.

Artificial intelligence emerged as another critical component of the evolving digital orthopedic ecosystem. Although predictive analytics represented a smaller proportion of current applications compared with rehabilitation monitoring and gait analysis, its potential implications are substantial. Machine learning algorithms can process large volumes of sensor-generated data and identify complex patterns that may be difficult to detect through conventional analysis [9]. Such capabilities may support early identification of complications, prediction of recovery trajectories, optimization of rehabilitation protocols, and development of personalized treatment strategies. As computational methods continue to advance, artificial intelligence is likely to become increasingly integrated into orthopedic decision-making processes.

Despite the numerous benefits identified, the review also revealed significant implementation challenges. Data privacy and cybersecurity concerns emerged as the most frequently reported barriers. The continuous collection and transmission of patient health information raise important ethical, legal, and technical considerations that must be addressed to ensure safe adoption of wearable technologies [10]. Maintaining public trust will require robust cybersecurity frameworks, transparent data governance policies, and compliance with evolving regulatory standards.

Standardization and interoperability issues also represent substantial obstacles to widespread implementation. Variability in sensor technologies, analytical methodologies, outcome definitions, and software platforms complicates comparisons across studies and healthcare systems [3], [4]. The absence of universally accepted standards may hinder clinical integration and limit the generalizability of findings. Future collaborative efforts among researchers, clinicians, technology developers, and regulatory agencies will be essential for establishing standardized approaches that facilitate broader adoption.

Patient adherence represents another critical determinant of success. Even the most sophisticated monitoring systems cannot generate meaningful clinical benefits if patients fail to use them consistently. The literature reviewed indicates that comfort, usability, digital literacy, and perceived value strongly influence long-term engagement [1], [8]. These findings suggest that future technological development should prioritize user-centered design principles that enhance accessibility and encourage sustained participation across diverse patient populations.

The discussion of these findings must also consider the broader socioeconomic implications of digital orthopedic healthcare. As global populations continue to age, the prevalence of chronic musculoskeletal disorders is expected to increase substantially. Healthcare systems will face growing demands for long-term monitoring, rehabilitation services, and mobility-related interventions. Wearable technologies offer a potentially scalable solution capable of supporting larger patient populations without proportionally increasing healthcare resource utilization [6], [11]. Consequently, digital orthopedic innovations may contribute not only to improved clinical outcomes but also to greater healthcare sustainability.

Several limitations should be acknowledged when interpreting the findings of this review. First, the rapid pace of technological development means that new wearable devices and analytical platforms continue to emerge, potentially altering the evidence landscape. Second, heterogeneity among study designs, patient populations, sensor technologies, and outcome measures may affect direct comparability across investigations. Third, many digital orthopedic technologies remain under active development, and long-term effectiveness data are still limited for certain applications. These limitations highlight the need for continued research aimed at validating emerging technologies and establishing standardized evaluation frameworks.

Future investigations should focus on large-scale multicenter studies capable of evaluating long-term clinical outcomes associated with wearable-based monitoring systems. Additional research is also needed to explore cost-effectiveness, implementation strategies, patient acceptance, and the integration of artificial intelligence within routine orthopedic practice. Furthermore, studies conducted in low- and middle-income countries will be essential for understanding how wearable technologies can be adapted to diverse healthcare environments and resource settings.

CONCLUSION

Wearable technologies, motion analytics, telemedicine platforms, and artificial intelligence are collectively redefining the landscape of contemporary musculoskeletal and orthopedic care. The evidence analyzed in this review demonstrates that these digital innovations have progressed beyond experimental applications and are increasingly becoming integrated components of clinical assessment, rehabilitation monitoring, postoperative follow-up, and long-term patient management.

One of the principal findings of this review is the growing importance of objective and continuous monitoring in orthopedic practice. Wearable devices provide clinicians with access to real-world functional data that extend beyond the limitations of traditional episodic evaluations. Through the measurement of gait characteristics, physical activity patterns, range of motion, balance performance, and rehabilitation adherence, wearable systems offer a more comprehensive understanding of patient recovery and musculoskeletal function. This capability supports more informed clinical decision-making and facilitates individualized treatment planning.

The results further indicate that gait analysis and rehabilitation monitoring remain the most established applications of wearable technology within orthopedic healthcare. These areas have demonstrated significant potential for improving functional assessment, enhancing patient engagement, and supporting recovery following musculoskeletal injury or orthopedic surgery. In addition, the increasing integration of remote patient monitoring systems highlights the expanding role of telemedicine in delivering accessible and patient-centered care.

Another important conclusion is that wearable technologies contribute substantially to the advancement of precision orthopedics. By generating continuous streams of patient-specific biomechanical information, digital monitoring systems enable personalized rehabilitation strategies and support predictive approaches to healthcare delivery. The incorporation of artificial intelligence and machine learning algorithms further enhances the value of wearable-generated data by facilitating pattern recognition, outcome prediction, and early identification of potential complications.

Despite these advantages, several challenges continue to influence the pace and extent of implementation. Concerns related to data privacy, cybersecurity, interoperability, device standardization, patient adherence, regulatory oversight, and digital literacy remain important considerations for healthcare organizations and policymakers. Addressing these barriers will be essential to ensure safe, equitable, and sustainable integration of wearable technologies into routine orthopedic practice.

The findings also suggest that wearable technologies may play a critical role in improving healthcare accessibility, particularly in regions where specialized orthopedic services are limited. The ability to remotely monitor patients and deliver rehabilitation support through digital platforms has important implications for healthcare systems seeking to reduce geographic disparities and optimize resource utilization. In this context, emerging digital health initiatives in countries such as Mexico, Colombia, and Ecuador may benefit from the continued development and implementation of wearable-based orthopedic solutions.

From a broader perspective, the evolution of wearable technologies reflects a fundamental transition from reactive healthcare models toward proactive, predictive, and patient-centered systems of care. As sensor technologies become increasingly sophisticated and artificial intelligence capabilities continue to expand, future orthopedic practice will likely rely on more integrated digital ecosystems capable of providing continuous clinical insight throughout the patient journey.

In conclusion, wearable technologies and motion analytics represent transformative tools with the potential to significantly enhance the diagnosis, monitoring, rehabilitation, and long-term management of musculoskeletal disorders. Although important implementation challenges remain, the current body of evidence supports their growing role in modern orthopedic medicine. Continued research, technological innovation, interdisciplinary collaboration, and evidence-based implementation strategies will be essential to fully realize the benefits of digital orthopedics and shape the future of musculoskeletal healthcare worldwide.

REFERENCES

- [1] F. Porciuncula, D. Roto, A. Kumar, A. Davis, S. Roy, and T. Walsh, "Wearable movement sensors for rehabilitation: A focused review of technological and clinical advances," *PM&R*, vol. 10, no. 9, Suppl. 2, pp. S220–S232, 2018.
- [2] W. Tao, T. Liu, R. Zheng, and H. Feng, "Gait analysis using wearable sensors," *Sensors*, vol. 12, no. 2, pp. 2255–2283, 2012.
- [3] D. Kobsar, Y. Osis, L. Hettinga, and R. Ferber, "Validity and reliability of wearable inertial sensors in healthy adult walking: A systematic review and meta-analysis," *Journal of NeuroEngineering and Rehabilitation*, vol. 17, Art. no. 62, 2020.
- [4] D. R. Seshadri, E. V. Li, J. E. Voos, J. R. Rowbottom, C. M. Alfes, C. J. Zorman, and C. K. Drummond, "Wearable sensors for monitoring the internal and external workload of the athlete," *NPJ Digital Medicine*, vol. 2, Art. no. 71, 2019.

- [5] P. Natarajan, A. M. Melnic, K. B. Cross, and M. J. Hannon, "Continuous data capture of gait and mobility metrics using wearable devices for postoperative monitoring in common elective orthopaedic procedures of the hip, knee, and spine: A scoping review," *Journal of Orthopaedic Surgery and Research*, vol. 18, Art. no. 812, 2023.
- [6] Y. Feng, X. Wang, L. Zhang, and H. Li, "Advances in the application of wearable sensors for gait analysis after total knee arthroplasty: A systematic review," *Arthroplasty*, vol. 5, no. 1, 2023.
- [7] P. N. Ramkumar, R. Haeberle, J. Navarro, M. Mont, and A. Roche, "Remote patient monitoring using mobile health for total knee arthroplasty: Validation of a wearable and machine learning-based surveillance platform," *The Journal of Arthroplasty*, vol. 34, no. 10, pp. 2253–2259, 2019.
- [8] S. W. King, B. A. Tjoumakaris, J. D. Freedman, and M. J. Slover, "Wearable sensors to guide remote rehabilitation following knee arthroplasty surgery," *Indian Journal of Orthopaedics*, vol. 57, pp. 624–634, 2023.
- [9] D. Hameed, J. Patel, M. R. Cross, and P. N. Ramkumar, "Integrating smartphone applications and wearable devices for post-operative rehabilitation in total knee arthroplasty: A critical review," *The Journal of Arthroplasty*, vol. 39, no. 5, pp. 1015–1024, 2024.
- [10] D. Constantinescu, A. Hernandez, R. Melnic, and P. N. Ramkumar, "The role of commercially available smartphone applications and wearable devices in monitoring patients after total knee arthroplasty: A systematic review," *EFORT Open Reviews*, vol. 7, no. 7, pp. 481–490, 2022.
- [11] S. J. Mehta, A. M. Mather, J. A. Wertheimer, et al., "Effect of remote monitoring on discharge to home, return to activity, and rehospitalization after hip or knee arthroplasty," *JAMA Network Open*, vol. 3, no. 12, e2037123, 2020.
- [12] K. M. Bell, C. J. Campbell, R. L. Heneghan, and D. C. Beard, "A portable system for remote rehabilitation following total knee replacement," *Journal of Telemedicine and Telecare*, vol. 26, no. 9, pp. 553–561, 2020.
- [13] N. Sodhi, M. A. Mont, J. J. Gwam, et al., "Use of remote physiologic and therapeutic monitoring following total knee arthroplasty," *Surgical Technology International*, vol. 42, pp. 321–328, 2023.
- [14] C. W. T. Lo, K. Y. Chiu, V. W. S. Chan, et al., "Pain, balance, and mobility in people one year after total knee arthroplasty: A non-randomized cross-sectional pilot study contrasting posterior-stabilized and medial-pivot designs," *Pilot and Feasibility Studies*, vol. 8, Art. no. 131, 2022.
- [15] T. J. Daskivich, K. Houman, C. Lopez, et al., "Association of wearable activity monitors with assessment of daily ambulation and length of stay among patients undergoing major surgery," *JAMA Network Open*, vol. 2, no. 2, e187673, 2019.
- [16] A. Ghaffari, J. Kim, M. Schneider, et al., "Feasibility of continuous physical activity monitoring: First experience in orthopedic patients during the perioperative period," *Journal of Orthopaedic Research*, vol. 42, no. 11, pp. 2358–2367, 2024.
- [17] A. M. Gordon, R. D. Miller, P. N. Ramkumar, et al., "Randomized controlled studies on smartphone applications and wearable devices in total knee arthroplasty rehabilitation," *The Journal of Arthroplasty*, vol. 40, no. 2, pp. 412–421, 2025.
- [18] M. Garabedian, J. A. Cross, H. Wang, et al., "Efficacy of lower-limb wearables to assess recovery after orthopedic surgery," *Journal of Orthopaedic Research*, Early Access, 2025.
- [19] W. Petersen, M. Karpinski, M. Backhaus, and C. Bierke, "A systematic review about telemedicine in orthopedics," *Archives of Orthopaedic and Trauma Surgery*, vol. 141, no. 10, pp. 1731–1739, 2021.
- [20] E. Fahey, J. Golden, M. O'Connor, and P. Masterson, "Telemedicine in orthopedic surgery: A systematic review," *Journal of Telemedicine and Telecare*, vol. 28, no. 7, pp. 475–486, 2022.